

Department of Surgery Division of Chest Surgery, Breast and Endocrinology Surgery

Toshiaki Morikawa, *Professor*
Tadashi Akiba, *Professor*
Shuji Sato, *Assistant Professor*
Makoto Odaka, *Assistant Professor*

Hiroshi Takeyama, *Professor*
Satoki Kinoshita, *Associate Professor*
Yasuo Toriumi, *Associate Professor*
Kazumi Kawase, *Assistant Professor*
Isao Tabei, *Assistant Professor*
Hiroko Nogi, *Assistant Professor*

General Summary

The Divisions of Chest Surgery, and of Breast and Endocrinology Surgery were established in June 2005. Since then, all staff members have been active in surgical practice, research, and education. Many studies are in progress.

Research Activities

Chest Surgery

Thoracoscopic surgery, is the focus of our clinical activity. This minimally invasive surgery produces fewer postoperative complications and sequelae and is especially beneficial for elderly, high-risk patients. Thoracoscopic surgery requires advanced skills, and we have independently developed total thoracoscopic surgery, which uses only a thoracoscope and video monitors to provide intraoperative views. Our method of thoracoscopic surgery can be used to treat many chest conditions, such as pneumothorax, peripheral lung nodules, mediastinal tumors, and lung cancer.

Thoracoscopic surgery is also indicated for higher-risk patients who have such complications as advanced pulmonary emphysema, impaired pulmonary function, and extremely high age and are not candidates for conventional open surgery.

Operative procedures, including wedge resection, segmentectomy, lobectomy, and pneumonectomy of the lung, are all safely performed, in addition to resection of mediastinal tumors or the thymus. Surgery for lung cancer requires much more advanced skills and oncological considerations, which have also been independently developed. Of the mediastinal procedures, thymectomy is usually performed via thoracoscopy rather than via a conventional median sternotomy. In our department the percentage of chest operations performed via thoracoscopy is more than 90%, which we assume to be the highest rate in the world.

The minimally invasive thoracoscopic surgery is being investigated with prospective clinical studies. These studies include a comparative study of open surgery and video-assisted surgery for lung cancer and evaluations of video-assisted surgery for bullous lung diseases in elderly patients with impaired lung function, of video-assisted surgery for thymic tumors, and of video-assisted thymectomy for myasthenia gravis.

Our clinical studies are also evaluating new devices and methods, such as narrow-band imaging for the thoracoscopic diagnosis of benign and malignant lung diseases, and Lapa-

roSonic coagulating shears (Ethicon Endo-Surgery, Inc., Cincinnati, OH, USA) for small thoracotomy. Three-dimensional diagnosis with computed tomography is used to make thoracoscopic surgery safer. The diagnosis and treatment of ground glass opacity of the lung, which is considered to indicate early adenocarcinoma, are being evaluated.

Many basic research studies are also underway. In the morphological expression-related advancement of the molecular genetic analysis of lung cancer, we are investigating whether CA19-9 activity is an important marker of de novo carcinogenesis. The biological and genetic characteristics of peripheral adenocarcinoma of the lung are being investigated to establish the most appropriate surgical procedures. The correlation of the detection of blood circulating tumor cells and the prognosis of patients with lung cancer is being examined.

The oncogenes of lung cancer are being analyzed with a next-generation sequencer.

A system for viewing videos on the Internet is now being developed and will help improve surgical training and research.

Breast

1. Clinical study

1) The evaluation of sentinel node biopsy after neoadjuvant chemotherapy

The minimally invasive technique of sentinel lymph node biopsy produces less morbidity and allows accurate pathologic staging of the axilla. Experience with sentinel node biopsy after neoadjuvant chemotherapy is limited. The purpose of our clinical study is to evaluate the feasibility, accuracy, and safety of this procedure in patients with breast cancer after neoadjuvant chemotherapy.

2) Evaluation of the usefulness of Sonazoid for detecting breast cancer

We performed phase II and III studies of the ultrasonographic imaging of the breast with the microbubble contrast medium Sonazoid (Daiichi Sankyo Co. Ltd., Tokyo) in collaboration with the Department of Radiology.

3) Evaluation of the effectiveness of exercise for psychiatric illness in patients after surgery for breast cancer

Many women experience a psychiatric illness, such as emotional distress, depression, and anxiety, after breast cancer is diagnosed. We have prospectively investigated the effects of exercise on psychological health.

4) Evaluation of the beneficial effect of cryotherapy for small cancers of the breast (Kashiwa Hospital)

5) Therapeutic strategy for oligometastatic breast cancer

We have analyzed patients with metastatic breast cancer for 30 years. The analysis indicates that oligometastatic breast cancer is a distinct subgroup with a long-term prognosis superior to that of metastatic breast cancer. We are performing prospective studies to better characterize oligometastatic breast cancer and to improve the prognosis of metastatic breast cancer.

2. Basic research

1) Development of breast cancer

We have used immunohistochemical techniques to investigate biological factors involved in the progression of carcinoma in situ to invasive breast cancer.

2) Clinically useful biomarkers for triple negative breast cancer

Triple-negative breast cancer is a heterogeneous disease. We have investigated prognostic and predictive biomarkers for triple-negative breast cancer.

3) Circulating tumor cells and disseminated tumor cells

Circulating tumor cells in the peripheral blood and disseminated tumor cells in the bone marrow of patients with breast cancer are strong prognostic factors. We have investigated their clinical values for patients with early breast cancers.

Endocrine

1. Basic research

1) Clinical usefulness of JT-95

A monoclonal antibody, designated JT-95, was made against a thyroid papillary carcinoma obtained by our Department of Breast and Endocrine Surgery. In collaboration with the Division of Molecular Cell Biology of The Jikei University, we have investigated the clinical usefulness of JT-95.

2) Association between JT-95 and cell-to-cell inhibition

We found that the adhesion between cells was inhibited in proportion to the quantity of JT-95 antibodies added in vitro. We have investigated the mechanism of cell-to-cell inhibition and association with lymph node metastases.

Publications

Nakada T, Akiba T, Inagaki T, Marushima H, Morikawa T, Ohki T. A case of a retroesophageal parathyroid adenoma with an aberrant right subclavian artery: a potential surgical pitfall. *Ann Thorac Cardiovasc Surg.* 2014; **20** Suppl: 786-9.

Nakada T, Akiba T, Inagaki T, Morikawa T, Ohki T. A rare case of primary intercostal leiomyoma: complete resection followed by reconstruction using a Gore-Tex® dual mesh. *Ann Thorac Cardiovasc Surg.* 2014; **20** Suppl: 617-21.

Akiba T, Nakada T, Inagaki T. A three-dimensional mediastinal model created with rapid prototyping in a patient with ectopic thymoma. *Ann Thorac Cardiovasc Surg.* 2015; **21**: 87-9.

Mimoto R, Kobayashi T, Imawari Y, Kamio M, Kato K, Nogi H, Toriumi Y, Hirooka S, Uchida K, Takeyama H. Clinical relevance and low tumor-initiating properties of oligometastatic breast cancer in pulmonary metastasectomy. *Breast Cancer Res Treat.* 2014; **147**: 317-24.

Akiba T, Morikawa T, Marushima H, Nakada T, Inagaki T, Ohki T. Computed tomography guided thoracoscopic segmentectomy for lung cancer with variant bronchus. *Ann Thorac Cardiovasc Surg.* 2014; **20**: 407-9.

Koido S, Kinoshita S, Mogami T, Kan S, Takakura K, Okamoto M, Homma S, Ohkusa T, Tajiri H, Harada J. Immunological assessment of cryotherapy in breast cancer patients. *Anticancer Res.* 2014; **34**: 4869-76.

Ito S, Araya J, Kurita Y, Kobayashi K, Taka-

saka N, Yoshida M, Hara H, Minagawa S, Wakui H, Fujii S, Kojima J, Shimizu K, Numata T, Kawaishi M, Odaka M, Morikawa T, Harada T, Nishimura SL, Kaneko Y, Nakayama K, Kuwano K. PARK2-mediated mitophagy is involved in regulation of HBEC senescence in COPD pathogenesis. *Autophagy.* 2015; **11**: 547-59.

Yamakawa H, Yoshida M, Yabe M, Baba Y, Baba E, Katagi H, Ishikawa T, Takagi M, Nakada T, Akiba T, Kuwano K. Pulmonary pleomorphic carcinoma detected as a result of pneumothorax and the subsequent occurrence of multiple cystic metastases. *Case Rep Med.* 2014; **2014**: 219273.

Nakada T, Akiba T, Inagaki T, Morikawa T, Ohki T. Simplified cavernostomy using wound protector for complex pulmonary aspergilloma. *Ann Thorac Surg.* 2014; **98**: 360-1.

Akiba T, Nakada T, Inagaki T. Simulation of the fissureless technique for thoracoscopic segmentectomy using rapid prototyping. *Ann Thorac Cardiovasc Surg.* 2015; **21**: 84-6.

Nakada T, Akiba T, Inagaki T, Morikawa T. Thoracoscopic anatomical subsegmentectomy of the right S2b + S3 using a 3D printing model with rapid prototyping. *Interact Cardiovasc Thorac Surg.* 2014; **19**: 696-8.

Akiba T, Inagaki T, Nakada T. Three-dimensional printing model of anomalous bronchi before surgery. *Ann Thorac Cardiovasc Surg.* 2014; **20**

Suppl: 659-62.

Akiba T, Nakada T, Inagaki T. Three-dimensional pulmonary model using rapid-prototyping in patient with lung cancer requiring segmentectomy. *Ann Thorac Cardiovasc Surg.* 2014; **20** Suppl: 490-2.

Kwong A, Chau WW, Kawase K. Work-life balance of female versus male surgeons in Hong Kong based on findings of a questionnaire designed by a Japanese surgeon. *Surg Today.* 2014; **44**: 62-72.