

Department of Anatomy (Gross Anatomy and Neuroanatomy)

Yoshinori Kawai, *Professor*

Tohru Hashimoto, *Assistant Professor*

General Summary

Our department research activities have focused on neuroanatomy and gross anatomy. In neuroanatomical research, organizations of neuronal networks and the development are investigated to elucidate brain function and diseases using morphological and electrophysiological methods. Our primary interest is focused on quantitative architecture and dynamics of neural circuits and their relationship. In gross anatomical researches, functional importance is explored on variations of organ systems using cadavers and animals.

Research Activities

To integrate and broadcast neural information, local microcircuits and global macrocircuits interact within certain specific nuclei of the central nervous system. The structural and functional architecture of this interaction was addressed for the caudal nucleus of the tractus solitarius (NTS), a relay station of peripheral viscerosensory information processed and conveyed to brain regions concerned with autonomic-affective and other interoceptive reflexive functions.

Geometric and functional architecture of viscerosensory microcircuitry

Is microcircuit wiring designed deterministically or probabilistically? Does geometric architecture predict functional dynamics of a given neuronal microcircuit? These questions were addressed in the viscerosensory microcircuit of the caudal nucleus of the tractus solitarius (NTS), which is generally thought to be homogeneous rather than laminar in cytoarchitecture. Using *in situ* hybridization histochemistry and whole-cell patch clamp recordings followed by neuronal reconstruction with biocytin filling, anatomical and functional organization of NTS microcircuitry was quantified to determine associative relationships. Morphologic and chemical features of NTS neurons displayed different patterns of process arborization and sub-nuclear localization according to neuronal types: smaller cells featured presynaptic local axons and GABAergic cells were aggregated specifically within the ventral NTS. The results suggested both a laminar organization and a spatial heterogeneity of NTS microcircuit connectivity. Geometric analysis of pre- and postsynaptic axodendritic arbor overlap of reconstructed neurons (according to parent somal distance) confirmed a heterogeneity of microcircuit connectivity that could underlie differential functional dynamics along the dorsoventral axis. Functional dynamics in terms of spontaneous and evoked postsynaptic current patterns behaved in a strongly location-specific manner according to the geometric dimension, suggesting a spatial laminar segregation of neuronal populations: a dorsal group of high excitation and a ventral

group of balanced excitation and inhibition. Recurrent polysynaptic activity was also noted in a subpopulation of the ventral group. Such geometric and functional laminar organization seems to provide the NTS microcircuit with both reverberation capability and a differentiated projection system for appropriate computation of visceral sensory information.