

## Erratum

### A Cross-sectional Survey on Smoking Cessation Counseling for Primary Care

Takuya NAGATA<sup>1,2</sup>, Masato MATSUSHIMA<sup>2</sup>, Tomokazu TOMINAGA<sup>2,3,5</sup>, Takamasa WATANABE<sup>4,5</sup>, and Yasuki FUJINUMA<sup>5,6</sup>

<sup>1</sup>*Ougibashi Clinic, Nankatsu Kinikyo*

<sup>2</sup>*Division of Clinical Epidemiology, The Jikei University School of Medicine*

<sup>3</sup>*Musashi-Koganei Clinic, Japanese Health and Welfare Co-operative Federation*

<sup>4</sup>*Kitaadachi Seikyo Clinic, Tokyo Hokuto Health Co-operative*

<sup>5</sup>*Centre for Family Medicine Development, Japanese Health and Welfare Co-operative Federation*

<sup>6</sup>*Interprofessional Education Research Center, Graduate School of Nursing, Chiba University*

The authors wish to apologize for confusion about the term, “primary-care smoking-related chronic disease (PC-SRCD)” in the paper, “A Cross-sectional Survey on Smoking Cessation Counseling for Primary Care : Jikeikai Med J. 2017 ; 64 : 23-30.”

The authors add the following sentences as a footnote ; “Primary-care smoking-related chronic disease (PC-SRCD)” is a term used in this article. There may be differences of opinion about chronic diseases related to smoking in primary care.

Furthermore, the authors wish to add the sentence, “The 95% confidence interval of kappa coefficient was calculated by bias-corrected bootstrap method.” in page 25, left column, line 33 to estimate more validly. As a result, the authors correct “95% confidence interval : 0.03-0.39” to “95% confidence interval : 0.02-0.45” in page 26, right column, line 7 and in page 23, line 10 of abstract. The authors think the conclusion of this research does not change.