

MS #25595

“Predictors associated with survival among elderly inpatients who receive cardiopulmonary resuscitation in Japan: an observational cohort study”

Tetsuro Hayashi, MD^{1,2}; Masato Matsushima, MD, MPH, PhD²; Seiji Bito, MD, MPH¹; Natsuko Kanazawa, MD, MPH^{3,4}; Norihiko Inoue, MD^{3,4}; Sarah Kyuragi Luthe, MD⁵; and Christina C. Wee, MD, MPH⁶

1) Division of Clinical Epidemiology, National Hospital Organization Tokyo Medical Center, Tokyo, Japan

2) Division of Clinical Epidemiology, The Jikei University School of Medicine, Tokyo, Japan

3) Department of Clinical Data Management and Research, Clinical Research Center, National Hospital Organization, Tokyo, Japan

4) Department of Health Policy and Informatics, Tokyo Medical and Dental University Graduate School, Tokyo, Japan

5) Department of Anesthesiology and Critical Care Medicine, Asahikawa Medical University, Hokkaido, Japan

6) General Medicine and Primary Care, Department of Medicine, Beth Israel Deaconess Medical Center, Boston, MA

Please address all correspondence to:

Tetsuro Hayashi, MD

Division of Clinical Epidemiology, National Hospital Organization Tokyo Medical Center, 2-5-1 Higashigaoka, Meguro-ku, Tokyo 152-8902, Japan

Tel: +81-3-3411-3130, Fax: +81-3-3411-3130

E-mail: tetsuro1026@hotmail.com

Coauthors' email address;

Masato Matsushima: masato@jikei.ac.jp

Seiji Bito: bitoseiji@kankakuki.go.jp

Natsuko Kanazawa: kanazawa-natsuko@hosp.go.jp

Norihiko Inoue: noricom@tmd.ac.jp

Sarah Kyuragi Luthe: sluthe@hsph.harvard.edu

Christina C. Wee: cwee@bidmc.harvard.edu

Running title: Cardiopulmonary resuscitation in elderly patients

Key words: aging, end of life care, decision making, evidence base medicine

Abstract word count (excluding sub-headings) = 292 words

Text word count (excluding abstract, acknowledgement, references, and tables) = 2117
words

Number of tables = 3

Number of figures = 0

Number of references = 22

Abstract

1 Background

2 Little is known about the outcomes of in-hospital cardiopulmonary resuscitation (CPR)
3 in Asian populations including elderly patients in Japan.

4 Objective

5 To determine the survival outcome of in-hospital CPR among elderly patients in Japan,
6 and to identify predictors associated with survival.

7 Design

8 Retrospective cohort study in 81 Japanese hospitals from April 1, 2010 to March 31,
9 2016.

10 Patients

11 We included elderly patients (age ≥ 65 years) who received CPR after two days of
12 hospitalization.

13 Main Measures

14 The primary outcome was survival at hospital discharge and the secondary outcomes
15 were the discharge disposition and consciousness level of patients who survived to
16 hospital discharge. To determine predictors associated with survival after in-hospital
17 CPR, we fit multivariable models for patient-level and institutional-level factors.

18 Key Results

19 Among the 5,365 patients who received CPR, 595 (11%) survived to discharge. Of
20 those who survived to discharge, 46% patients were discharged home, and 10% patients
21 were comatose at discharge. Older age and higher burden of comorbidities were
22 associated with reduced survival. The adjusted OR was 0.35 (95% CI, 0.22-0.55) for
23 age >90 years compared to age 65-69 years, and 0.68 (95% CI, 0.48-0.97) for Charlson
24 Comorbidity Index score of ≥ 4 compared with score of 0. Other predictors of reduced
25 survival included receiving CPR on weekends compared to weekdays (AOR, 0.63; 95%

26 CI, 0.51-0.77) and in small hospitals compared to large hospitals (AOR, 0.58; 95% CI,
27 0.40-0.83).

28

29 **Conclusions**

30 Among elderly patients in Japan, the survival rate of in-hospital CPR was
31 approximately one in ten, and less than half of these patients were discharged home. In
32 addition to older age and higher illness burden, receiving CPR on weekends and/or in
33 small hospitals were significant predictors of reduced survival. These findings should be
34 considered in advanced care planning discussions with elderly patients to avoid
35 subjecting patients to CPR that are likely futile.

36 **Introduction**

37 Whether patients should receive cardiopulmonary resuscitation (CPR) is an important
38 decision encountered by elderly patients and clinical teams. However, outcome data on
39 inpatient CPR is unclear. Additionally, it has been reported that elderly patients and
40 physicians overestimate the chance of survival after CPR when deciding whether a
41 do-not-resuscitate (DNR) order might be appropriate.¹⁻⁵ Therefore, providing accurate
42 outcomes on CPR may influence the decision making on CPR for hospitalized elderly
43 patients.⁶

44

45 A recent systematic review including 29 studies showed that the survival rate of CPR
46 among elderly inpatients ranges from 11.6% to 18.7% and declines with increasing age.
47 The vast majority of these patients were from the US,⁷ and survival rate after CPR has
48 also been shown to vary by race.⁸ A previous study using Medicare data reported that
49 black and other non-white patients had a higher likelihood of receiving in-hospital CPR
50 but had lower odds of survival.⁹

51

52 In contrast, only a few studies have examined the outcomes of in-hospital CPR in Asian
53 populations including elderly patients in Japan, who have the longest life expectancy in
54 the world.¹⁰ In Japan, more than 70% of hospitalized patients are aged ≥ 65 years,
55 approximately three in four die in the hospital, and two in ten do not have DNR
56 orders.¹¹ In this context, we used a large population-based database from 81 Japanese
57 hospitals to investigate the survival rate after CPR among hospitalized elderly patients
58 and to identify predictors associated with survival.

59

60 **Methods**

61 *Date source*

62 Patients' data were extracted from the Diagnosis Procedure Combination (DPC) data¹²
63 of National Hospital Organization (NHO) Network in Japan. The DPC is nationally
64 used for health care insurance claims for health service rendered in acute care hospitals
65 in Japan, similar to medical claims codes used in the US.^{13,14} The NHO is the largest
66 hospital network in Japan, and stores DPC data from 81 acute care hospitals, typically
67 teaching/tertiary hospitals in each district, affiliated with the NHO for administration
68 and clinical information analysis. The DPC data includes hospital administrative data
69 and discharge abstracts: unique identifiers of hospitals; characteristics of patients;
70 admission and discharge status; diagnoses and comorbidities at admission, and
71 complications after admission recorded in the International Classification of Diseases,
72 Tenth Revision (ICD-10); surgical and non-surgical procedures; drugs and devices used;
73 and length of stay.¹³ To optimize the accuracy of medical information, attending
74 physicians are responsible for registering the diagnoses. This retrospective study was
75 approved by the Institutional Ethics Committees at the National Hospital Organization
76 and National Tokyo Medical Center.

77

78 *Study sample*

79 This study included patients aged ≥ 65 years who were hospitalized and received CPR
80 two days after admission from April 1, 2010 to March 31, 2016. We excluded patients
81 who received CPR within two days of hospitalization due to the inability to distinguish
82 patients who were in cardiac arrest on arrival to the hospitals. The procedure of CPR
83 was identified as "Closed-chest cardiac massage (J046)" from the DPC database. For
84 patients who went into more than one cardiac arrest event and received multiple CPR
85 episodes during their hospitalization, we only included the first CPR in our analysis.

86

87 *Outcome variables*

88 Our primary outcome was survival to hospital discharge among elderly inpatients who
89 received CPR after the second day of their hospitalization. As secondary outcomes, we
90 investigated the discharge disposition (home, nursing facilities, or other hospitals) of
91 patients who survived to hospital discharge and the patient's level of consciousness at
92 discharge (comatose or not). Consciousness level was categorized based on the Japan
93 Coma Scale where comatose is defined by the 3-digit code of 100, 200 or 300.¹⁵

94

95 *Potential predictors*

96 We considered the following potential patient-level and institutional-level predictors:
97 patient sex, age, body mass index (BMI), consciousness level on admission (based on
98 the Japan Coma Scale),¹⁵ admitting diagnosis, admitting comorbidities (based on the
99 Charlson comorbidity index),¹⁶ hospital size (defined by number of hospital beds: <300,
100 small; 300-499, medium; and ≥ 500 , large), days from admission to CPR, and day of
101 week of CPR (weekend vs. weekday). We also considered whether patients received the
102 following interventions within three days prior to CPR in order to assess
103 pre-resuscitation interventions: intensive care unit (ICU) admission, enteral nutrition,
104 total parenteral nutrition, vasopressor use, and mechanical ventilation (defined by
105 non-invasive positive pressure ventilation and/or invasive mechanical ventilation).

106

107 *Statistical methods*

108 We compared baseline patient characteristics using chi-square test. To determine the
109 predictors associated with survival after in-hospital CPR at the level of $\alpha = 0.05$, we
110 fit unadjusted and adjusted logistic regression models for each response using a forward
111 selection approach. Non-significant variables were added sequentially; no potential
112 confounders that altered estimates of significant factors by more than 10% were
113 identified. Thus, candidate explanatory variables associated with survival were
114 identified. Next, we analyzed the candidate variables using generalized estimating

115 equations with a logit link function where the clustering effect associated with hospitals
116 is accounted for by the robust sandwich standard error estimator. We tested the
117 significance of BMI in a subset of patients with complete data as BMI was missing in
118 885 patients (16%), and found no significant importance. We also performed a
119 sensitivity analysis excluding patients who underwent multiple CPR events. All
120 analyses were performed with STATA 12 software (STATA Corp, College Station, TX).
121 All P values were two-tailed and considered statistically significant with $P < 0.05$.

122

123 **Results**

124 *Patients*

125 We identified 1,478,934 patients aged ≥ 65 years who were hospitalized for more than
126 two days in 81 Japanese hospitals from April 1, 2010 to March 31, 2016. Among these,
127 78,360 patients experienced cardiac arrest after two days of admission, and 5,365
128 patients received CPR.

129

130 *Descriptive statistics*

131 Among the 5,365 patients who received CPR, 595 patients (11%) survived to hospital
132 discharge. Of the 4,770 patients who died after CPR, 3,768 patients (79%) died within
133 the same day of receiving CPR. Overall, the median length of stay was 22 days, and the
134 median length from hospital admission to event of cardiac arrest was 16 days. Table 1
135 shows the characteristics of the patients who received CPR and the survival rates. The
136 median age was 79 years (IQR, 73-84 years) and 63% were male. Of 231 patients who
137 underwent multiple CPR events during hospitalization, 32 (14%) patients survived to
138 hospital discharge.

139

140 *Status among patients who survived to discharge after CPR*

141 Among 595 patients who survived to discharge, the median length of hospitalization

142 from receiving CPR to discharge was 46 days, and 59 (10%) patients were comatose at
143 discharge (Table 2). Of 581 patients with data available on discharge dispositions, 279
144 (48%) patients were transferred to other hospitals, 266 (46%) patients were discharged
145 home, and 18 (3%) patients were transferred to nursing facilities.

146

147 *Multivariable analysis*

148 Table 3 presents significant predictors associated with survival to hospital discharge
149 after adjustment. Age and higher burden of comorbidities were significant predictors of
150 reduced survival. Admitting diagnoses of cancer, infectious disease, and hematological
151 disease were also significant predictors of reduced survival, while admitting diagnoses
152 of ischemic heart disease, arrhythmia on admission, and ICU admission prior to
153 resuscitation were associated with greater survival. The patients who received CPR after
154 two weeks from admission were less likely to survive. Receiving CPR on weekends
155 and/or in small hospitals were also significant predictors of reduced survival. There was
156 no effect modification between hospital size and receiving CPR on weekends or
157 between hospital size and ICU admission. Other factors such as sex, BMI,
158 consciousness level on admission, and pre-resuscitation interventions of enteral
159 nutrition, total parental nutrition, vasopressor treatment, and mechanical ventilation
160 were not significant predictors or confounders. In the sensitivity analysis excluding
161 patients who underwent multiple CPR events, the significance and direction of the
162 effect of the predictors were not different from the result in the main analysis.

163

164

165 **Discussion**

166 In our study of over 5,000 elderly patients who received in-hospital CPR in Japan,
167 approximately one in ten survived, and less than half of these patients were discharged
168 home. In addition to older age and higher illness burden, receiving CPR on weekends
169 and/or in small hospitals were significant predictors of reduced survival; in contrast,
170 ischemic heart disease, arrhythmia on admission, and ICU admission before
171 resuscitation were associated with greater survival.

172

173 To the best of our knowledge, this is the largest study outside of the US examining the
174 outcomes of in-hospital CPR among elderly patients, and the first study focusing on an
175 Asian population. Further, this study considered pre-resuscitation interventions and
176 institutional-level factors in addition to baseline characteristics on hospital admission.

177

178 Approximately half of the patients who survived to discharge after CPR were
179 discharged home. In contrast, another half of those were transferred to other hospitals;
180 one in ten were comatose at discharge. Many of the patients transferred to other
181 hospitals were likely to move to rehabilitation hospitals or long-term care hospitals
182 rather than other acute care hospitals though these data were not available in our study.
183 Nonetheless, these findings suggest that these patients had physical complications after
184 CPR and required longer-term medical treatment.

185

186 Our findings are consistent with prior data reporting the association between older age
187 and reduced survival.^{7,9} A systematic review showed that the pooled survival rate after
188 CPR was 18.7% for patients between 70 and 79 years old, 15.4% for those between 80
189 and 89 years old, and 11.6% for 90 years and older.⁷ The survival rates of in-hospital
190 CPR in the present study were even lower than that of prior studies. This difference may
191 be partly explained by the different inclusion criteria of when CPR was performed; for
192 example, we excluded patients who received CPR within two days of hospitalization
193 due to our inability to distinguish patients who were in cardiac arrest on arrival to the
194 hospitals.

195

196 Our finding that arrhythmia, ischemic heart disease, and ICU admission were associated
197 with increased survival rates likely reflects increased use of cardiac monitoring and
198 rapid response to event.¹⁷ In addition, patients with arrhythmia or ischemic heart disease

199 might be more likely to have initial rhythms of ventricular fibrillation or ventricular
200 tachycardia, which are more responsive to CPR.^{18,19} Furthermore, there may be a
201 selection bias in which patients are admitted to ICU based on physician's perception of
202 patient's likelihood to benefit from ICU care. In contrast, cancer, infection,
203 hematological disease and higher burden of comorbidities were independently
204 predictive of in-hospital mortality, and these findings are in line with previous
205 studies.^{9,17} These predictors of poor prognosis can be applied to advanced care planning
206 discussions and guide decision-making about do-not-resuscitate orders.

207

208 Our finding that receiving CPR on weekends compared to weekdays as an independent
209 predictor of reduced survival is consistent with a previous study demonstrating that
210 patients who received CPR during day on weekdays were more likely to survive,
211 compared to those who received CPR during day on weekends.²⁰ Our findings also
212 indicate that receiving CPR in smaller hospitals compared to larger hospitals were
213 associated with reduced survival. The underlying mechanism of reduced survival after
214 CPR on weekends and in small hospitals may be explained by different hospital staffing
215 patterns in these settings. The difference in the availability of rapid response systems
216 (RRS) between larger hospitals and smaller hospitals may be another explanation, since
217 the shortage of medical staff is considered to be the main barrier of implementing
218 RRS.²¹ Addressing institutional systems, such as implementing RRS among small
219 hospitals during weekends, may improve important outcomes associated with CPR.

220

221 While our study leverages a rich dataset, it is not without limitations. First, our
222 definition of CPR was based on the DPC data in Japan, and the definition of CPR has
223 not been validated in this data, although a previous validation study on the DPC data
224 shows that the sensitivity and specificity of common procedures exceeds 90% and the
225 sensitivity and specificity of primary diagnoses were 79% and 93%.²² Second, our

226 analyses were limited to patients who received CPR after two days of hospitalization,
227 which may have underestimated the survival rate in our study. Third, the dataset had
228 limited information on other potential predictors of survival such as initial arrest rhythm,
229 presence of witness, use of telemetry, and the availability of rapid response systems. In
230 addition, the dataset had also limited information on other outcomes such as physical
231 function, quality of life, and details of hospitals to which patients were transferred.

232

233 In conclusion, among elderly patients in Japan, the survival rate of in-hospital CPR after
234 two days of hospitalization was approximately one in ten, and less than half of these
235 patients were discharged home. In addition to older patient age and higher illness
236 burden, receiving CPR on weekends and in small hospitals were significant predictors
237 of reduced survival. These findings suggest that patients' baseline status should be
238 considered in advanced care planning discussions with elderly patients to avoid
239 subjecting patients to CPR that are likely futile. Moreover, future studies should
240 identify potential system factors that might underlie differences in outcomes after CPR
241 between large and small hospitals.

242

243

244 **Acknowledgments**

245 *Contributors*

246 We thank Mr. Masaya Nakadera for the assistance with data extraction and thank Dr.
247 Suminobu Ito and Dr. Hiromasa Horiguchi for organizing this research.

248

249 *Funding*

250 This study was supported by a research grant for post-graduate students from Jikei
251 University School of Medicine. The study sponsor had no role in the study design, data
252 collection, analysis and interpretation, report writing, or the decision to submit the

253 article for publication. MM is the program director of Jikei Clinical Research Program
254 for Primary Care, and TH was a former trainee of the Jikei Clinical Research Program
255 for Primary Care. When this study was conducted, TH was the Shigeaki Hinohara, MD,
256 International Primary Care Fellow at Beth Israel Deaconess Medical Center (BIDMC)
257 which was supported by the Ryoichi Sasakawa Fellowship Fund at BIDMC. SL was
258 supported by grants from World Bank and Kamiyama Foundation. CW is supported by
259 Midcareer Mentorship Award from the National Institutes of Health (K24DK087932).

260

261 *Presentation*

262 We presented an earlier version of the manuscript as a poster at the 2018 SGIM annual
263 meeting on April 12th, 2018.

264

265 **Conflict of interest**

266 The authors declare that they do not have a conflict of interest.

267

268 **References**

- 269 1. Miller DL, Jahnigen DW, Gorbien MJ, Simbartl L. Cardiopulmonary
270 resuscitation: how useful? Attitudes and knowledge of an elderly population.
271 *Archives of internal medicine*. 1992;152(3):578-582.
- 272 2. Schonwetter RS, Walker RM, Kramer DR, Robinson BE. Resuscitation decision
273 making in the elderly: the value of outcome data. *Journal of general internal*
274 *medicine*. 1993;8(6):295-300.
- 275 3. Mead GE, Turnbull CJ. Cardiopulmonary resuscitation in the elderly: patients'
276 and relatives' views. *Journal of medical ethics*. 1995;21(1):39-44.
- 277 4. Frank C, Heyland DK, Chen B, Farquhar D, Myers K, Iwaasa K. Determining
278 resuscitation preferences of elderly inpatients: a review of the literature. *CMAJ :*
279 *Canadian Medical Association journal = journal de l'Association medicale*
280 *canadienne*. 2003;169(8):795-799.
- 281 5. Miller DL, Gorbien MJ, Simbartl LA, Jahnigen DW. Factors influencing
282 physicians in recommending in-hospital cardiopulmonary resuscitation. *Archives*
283 *of internal medicine*. 1993;153(17):1999-2003.
- 284 6. Murphy DJ, Burrows D, Santilli S, et al. The influence of the probability of
285 survival on patients' preferences regarding cardiopulmonary resuscitation. *The*
286 *New England journal of medicine*. 1994;330(8):545-549.
- 287 7. van Gijn MS, Frijns D, van de Glind EM, B CvM, Hamaker ME. The chance of
288 survival and the functional outcome after in-hospital cardiopulmonary
289 resuscitation in older people: a systematic review. *Age and ageing*.
290 2014;43(4):456-463.
- 291 8. Galea S, Blaney S, Nandi A, et al. Explaining racial disparities in incidence of
292 and survival from out-of-hospital cardiac arrest. *American journal of*
293 *epidemiology*. 2007;166(5):534-543.
- 294 9. Ehlenbach WJ, Barnato AE, Curtis JR, et al. Epidemiologic study of in-hospital
295 cardiopulmonary resuscitation in the elderly. *The New England journal of*
296 *medicine*. 2009;361(1):22-31.
- 297 10. Tokudome S, Hashimoto S, Igata A. Life expectancy and healthy life expectancy
298 of Japan: the fastest graying society in the world. *BMC research notes*.
299 2016;9(1):482.

- 300 11. Sasaki A, Hiraoka E, Homma Y, et al. Association of code status discussion with
301 invasive procedures among advanced-stage cancer and noncancer patients.
302 *International journal of general medicine*. 2017;10:207-214.
- 303 12. Tanaka S, Seto K, Kawakami K. Pharmacoepidemiology in Japan: medical
304 databases and research achievements. *Journal of pharmaceutical health care*
305 *and sciences*. 2015;1:16.
- 306 13. Sumitani M, Uchida K, Yasunaga H, et al. Prevalence of malignant hyperthermia
307 and relationship with anesthetics in Japan: data from the diagnosis procedure
308 combination database. *Anesthesiology*. 2011;114(1):84-90.
- 309 14. Isogai T, Yasunaga H, Matsui H, et al. Effect of weekend admission for acute
310 myocardial infarction on in-hospital mortality: a retrospective cohort study.
311 *International journal of cardiology*. 2015;179:315-320.
- 312 15. Shigematsu K, Nakano H, Watanabe Y. The eye response test alone is sufficient
313 to predict stroke outcome--reintroduction of Japan Coma Scale: a cohort study.
314 *BMJ open*. 2013;3(4).
- 315 16. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying
316 prognostic comorbidity in longitudinal studies: development and validation.
317 *Journal of chronic diseases*. 1987;40(5):373-383.
- 318 17. Larkin GL, Copes WS, Nathanson BH, Kaye W. Pre-resuscitation factors
319 associated with mortality in 49,130 cases of in-hospital cardiac arrest: a report
320 from the National Registry for Cardiopulmonary Resuscitation. *Resuscitation*.
321 2010;81(3):302-311.
- 322 18. Nadkarni VM, Larkin GL, Peberdy MA, et al. First documented rhythm and
323 clinical outcome from in-hospital cardiac arrest among children and adults.
324 *Jama*. 2006;295(1):50-57.
- 325 19. Peberdy MA, Kaye W, Ornato JP, et al. Cardiopulmonary resuscitation of adults
326 in the hospital: a report of 14720 cardiac arrests from the National Registry of
327 Cardiopulmonary Resuscitation. *Resuscitation*. 2003;58(3):297-308.
- 328 20. Peberdy MA, Ornato JP, Larkin GL, et al. Survival from in-hospital cardiac
329 arrest during nights and weekends. *Jama*. 2008;299(7):785-792.
- 330 21. Koike T, Nakagawa M, Shimosawa N, et al. The implementation of rapid
331 response system in Japanese hospitals: its obstacles and possible solutions.

- 332 *Journal of Japanese association for acute medicine*. 2017;28(6):219-229 (in
333 Japanese).
- 334 22. Yamana H, Moriwaki M, Horiguchi H, Kodan M, Fushimi K, Yasunaga H.
335 Validity of diagnoses, procedures, and laboratory data in Japanese administrative
336 data. *Journal of epidemiology*. 2017;27(10):476-482.
- 337

338 **Table 1.** Characteristics of patients who received in-hospital cardiopulmonary resuscitation (N=5,365)

339

Characteristics	Sample		Survival rate	P-value*
	n	(%)	%	
Overall	5,365	(100.0)	11.1	
Sex				
Male	3,375	(62.9)	10.8	0.35
Female	1,990	(37.1)	11.6	
Age				
65-69	677	(12.6)	13.6	0.006
70-74	890	(16.6)	9.9	
75-79	1,181	(22.0)	12.1	
80-84	1,308	(24.4)	11.1	
85-89	888	(16.6)	11.3	
≥90	421	(7.9)	6.4	
Body Mass Index†				
<18.5	1,251	(23.3)	9.4	0.06
18.5-25	3,487	(65.0)	11.4	
≥25	627	(11.7)	12.8	
Comatose on admission	296	(5.5)	15.2	0.02
Admitting diagnosis				
Cancer	984	(18.3)	6.5	< 0.001
Pneumonia	543	(10.1)	9.8	
Congestive heart failure	542	(10.1)	13.5	
Cerebrovascular diseases	377	(7.0)	13.0	
Other respiratory diseases	316	(5.9)	7.9	
Traumatic diseases	291	(5.4)	14.8	
Ischemic heart diseases	290	(5.4)	21.7	
Gastroenterological diseases	183	(3.4)	12.0	
Other infectious diseases‡	168	(3.1)	4.8	
Aortic dissection	122	(2.3)	13.1	
Neurological diseases	101	(1.9)	13.9	
Renal failure	98	(1.8)	12.2	
Hematological diseases	92	(1.7)	1.1	
Arrhythmia	88	(1.6)	33.0	

Other	1,170 (21.8)	10.5	
Charlson comorbidity index			
0	1,669 (31.1)	12.2	
1	1,415 (26.4)	12.4	
2	1,016 (18.9)	10.5	0.009
3	714 (13.3)	8.8	
≥4	551 (10.3)	8.2	
Pre-resuscitation interventions			
Intensive care unit admission	532 (9.9)	17.9	< 0.001
Enteral nutrition	820 (15.3)	11.2	0.90
Total parental nutrition	758 (14.1)	8.8	0.03
Vasopressor use	1,103 (20.6)	12.1	0.21
Mechanical ventilation	854 (15.9)	13.3	0.02
Days from admission to CPR			
			< 0.001
3-7	1,383 (25.8)	14.3	
8-14	1,064 (19.8)	12.3	
15-28	1,174 (21.9)	9.5	
≥29	1,744 (32.5)	8.8	
Day of CPR			
Weekday	3,695 (68.9)	12.4	
Weekend	1,670 (31.1)	8.3	< 0.001
Hospital size (number of hospital beds)			
<300	845 (15.8)	6.2	
300-499	3,094 (57.7)	12.0	< 0.001
≥500	1,426 (26.6)	12.1	

340 Abbreviation: CPR, cardiopulmonary resuscitation

341 * All the comparisons were made using chi-square tests; † 16 % missing; ‡ including sepsis

342 **Table 2.** Status among patients who survived to discharge after cardiopulmonary resuscitation (N=595)

343

	n (%)
Discharge disposition*	
Other hospitals	279 (48.0)
Home	266 (45.8)
Nursing facilities	18 (3.1)
Other	18 (3.1)
Neurological status	
Comatose at discharge	59 (9.9)

344

* 581 patients' data (98%) were available on discharge dispositions

345 **Table 3.** Multivariable analyses for predictors associated with survival to hospital discharge*

Predictors	Odds Ratio (95%CI)
Age	
65-69	Reference
70-74	0.69 (0.50-0.95)
75-79	0.84 (0.63-1.12)
80-84	0.7 (0.52-0.94)
85-89	0.71 (0.52-0.97)
≥90	0.35 (0.22-0.55)
Admitting diagnosis	
Pneumonia	Reference
Cancer	0.61 (0.41-0.90)
Congestive heart failure	1.32 (0.9-1.94)
Cerebrovascular diseases	1.16 (0.76-1.76)
Other respiratory diseases	0.73 (0.44-1.21)
Traumatic diseases	1.46 (0.94-2.26)
Ischemic heart diseases	1.84 (1.21-2.80)
Gastroenterological diseases	1.19 (0.69-2.03)
Other infectious diseases†	0.4 (0.19-0.87)
Aortic dissection	1.04 (0.56-1.92)
Neurological diseases	1.36 (0.72-2.59)
Renal failure	1.14 (0.58-2.25)
Hematological diseases	0.08 (0.01-0.63)
Arrhythmia	3.96 (2.30-6.79)
Other	1.02 (0.72-1.44)
Charlson comorbidity index	
0	Reference
1	0.99 (0.79-1.24)
2	0.83 (0.64-1.07)
3	0.75 (0.55-1.03)
≥4	0.68 (0.48-0.97)
Pre-resuscitation intervention	
Intensive care unit admission	1.38 (1.06-1.80)
Days from admission to CPR	
3-7	Reference
8-14	0.88 (0.69-1.12)

15-28	0.68 (0.53-0.88)
≥29	0.67 (0.53-0.85)
Day of CPR	
Weekday	Reference
Weekend	0.63 (0.51-0.77)
Number of hospital beds	
<300	0.58 (0.40-0.83)
300-499	1.05 (0.83-1.32)
≥500	Reference

346 Abbreviation: CPR, cardiopulmonary resuscitation

347 * Adjusted for the year of admission; † including sepsis

1 **Supplementary appendix**

2 **Table S1.** Sensitivity analysis for predictors associated with survival to hospital discharge excluding
 3 patients who underwent multiple CPR events (N=5,134) *

4

Predictors	Odds Ratio (95%CI)
Age	
65-69	Reference
70-74	0.72 (0.51-0.99)
75-79	0.83 (0.61-1.12)
80-84	0.72 (0.54-0.98)
85-89	0.72 (0.52-1.00)
≥90	0.34 (0.21-0.54)
Admitting diagnosis	
Pneumonia	Reference
Cancer	0.62 (0.42-0.92)
Congestive heart failure	1.27 (0.86-1.89)
Cerebrovascular diseases	1.12 (0.73-1.72)
Other respiratory diseases	0.74 (0.45-1.23)
Traumatic diseases	1.45 (0.94-2.26)
Ischemic heart diseases	1.71 (1.10-2.65)
Gastroenterological diseases	1.18 (0.69-2.02)
Other infectious diseases†	0.31 (0.13-0.74)
Aortic dissection	1.09 (0.59-2.02)
Neurological diseases	1.33 (0.68-2.57)
Renal failure	1.19 (0.60-2.35)
Hematological diseases	0.09 (0.01-0.64)
Arrhythmia	3.94 (2.27-6.83)
Other	0.99 (0.70-1.40)
Charlson comorbidity index	
0	Reference
1	0.97 (0.78-1.22)
2	0.83 (0.64-1.07)
3	0.75 (0.54-1.03)
≥4	0.69 (0.48-0.98)
Pre-resuscitation intervention	

Intensive care unit admission	1.35 (1.02-1.79)
Days from admission to CPR	
3-7	Reference
8-14	0.89 (0.69-1.15)
15-28	0.72 (0.56-0.94)
≥29	0.69 (0.55-0.88)
Day of CPR	
Weekday	Reference
Weekend	0.63 (0.51-0.77)
Number of hospital beds	
<300	0.58 (0.40-0.83)
300-499	1.04 (0.82-1.31)
≥500	Reference

5 Abbreviation: CPR, cardiopulmonary resuscitation

6 * Adjusted for the year of admission; † including sepsis

7