

Institute of DNA Medicine

Department of Molecular Immunology

Saburo Saito, Associate Professor and Director
Nobutake Akiyama, Assistant Professor

Daitaro Kurosaka, Associate Professor
Yuji Ohno, Assistant Professor

General Summary

Our research interests have focused mainly on analysis of the basic immune system to protect against diseases and of immune disorders, such as hypersensitivity diseases and autoimmune diseases.

Research Activities

Pleiotropic function of interleukin-31

Interleukin (IL)-31 is a T-cell-derived cytokine that induces severe pruritus, hair loss, and dermatitis and is involved in allergic diseases, such as atopic dermatitis and bronchitis. To investigate the function of IL-31, IL-31 transgenic (Tg) mice were generated in our laboratory. In addition to scratching behavior and hair loss as reported previously, enhancement of serum immunoglobulin (Ig) E level was observed in the IL-31 Tg mice. Furthermore, these pleiotropic functions were verified by the administration of IL-31 into normal mice. To further analyze the mechanism of IgE production by IL-31, we are seeking factors enhancing T helper type 2 (Th2) cytokine production, focusing on the IL-31 receptor-expressing cells, such as keratinocytes, macrophages, and granulocytes.

A rice-based edible vaccine expressing Japanese cedar pollen allergens induces oral tolerance in Japanese monkeys with Japanese cedar pollinosis

Japanese cedar (*Cryptomeria japonica*: CJ) pollinosis affects more than 30% of the Japanese population and is, thus, one of the most common diseases in Japan. Furthermore, CJ pollinosis has been found to occur naturally in Japanese monkeys (*Macaca fuscata*), which show symptoms similar to those of human patients.

Plants have recently been recognized as a form of bioreactor for the cost-effective production of large-scale recombinant proteins. The edible tissue of plants further provides the significant benefit of achieving a simple method of mucosal delivery of vaccines without the need for complicated purification steps.

Our previous study showed that oral administration of Tg rice seeds that have accumulated high concentrations of polypeptides derived from CJ pollen allergens to mice reduces their serum IgE levels and T-cell proliferative responses to CJ allergens, proving the efficacy of oral immunotherapy for the treatment of pollinosis.

In this study, the Tg rice plants that had accumulated high concentrations of JC allergens were used for oral immunotherapy for CJ pollinosis of monkeys. Five monkeys with CJ pollinosis were fed once a day with 20 g of the rice seeds containing about 50 to 60 mg of

allergens for 3 months. No side effects, such as urticaria, dyspnea, vomiting, and weight loss, were observed during immunotherapy. One and a half months after the start of feeding, proliferative responses of T cells to JC allergens in 4 of 5 monkeys were significantly inhibited compared with those in monkeys at the start of feeding. However, their T-cell responses to CJ allergens were restored 1 month after the end of feeding. These results indicate that oral immunotherapy with Tg rice seeds is safe and effective for the treatment of pollinosis.

Construction of a new anticancer strategy focused on glycosylation

We are developing a novel anticancer strategy that induces cytotoxic T cells against non-polarized cells represented by cancer cells, by enhancing MHC class I-restricted antigen presentation by inhibiting N-glycosylation.

Analysis of the N-glycosylation structure that controls the secretion of IL-31 showed that some structures of N-glycosylation were able to enhance MHC class I-restricted antigen presentation. On the basis of this finding, we are developing a new vaccine that induces cytotoxic T cells against cancer or viruses with artificial immature N-glycosylated proteins.

Publications

Fukuda T, Akiyama N, Ikegami M, Takahashi H, Sasaki A, Oka H, Komori T, Tanaka Y, Nakazato Y, Akimoto J, Tanaka M, Okada Y, Saito S. Expression of hydroxyindole-o-methyltransferase enzyme in the human central nervous system and in pineal parenchymal cell tumors. *J Neuropathol Exp Neurol* 2010; **69**: 498-510.

Kurosaka D, Hirai K, Nishioka M, Miyamoto Y, Yoshida K, Noda K, Ukichi T, Yanagimachi M, Furuya K, Takahashi E, Kingetsu I, Fukuda K, Yamada A. Clinical significance of serum levels

of vascular endothelial growth factor, angiopoietin-1, and angiopoietin-2 in patients with rheumatoid arthritis. *J Rheumatol* 2010; **37**: 1121-8.

Yoshida K, Kurosaka D, Joh K, Matsushima S, Takahashi E, Hirai K, Noda K, Ukichi T, Furuya K, Yanagimachi M, Kingetsu I, Fukuda K, Yamada A. Fasciitis as a common lesion of dermatomyositis, demonstrated early after disease onset by en bloc biopsy combined with magnetic resonance imaging. *Arthritis Rheum* 2010; **62**: 3751-9.