

Department of Internal Medicine

Division of Diabetes, Metabolism and Endocrinology

Kazunori Utsunomiya, *Professor*
Junichi Yokoyama, *Professor*
Katsuyoshi Tojo, *Professor*
Yutaka Mori, *Professor*
Masami Nemoto, *Associate Professor*
Rimei Nishimura, *Associate Professor*

Yoichi Sakamoto, *Professor*
Takashi Sasaki, *Professor*
Kuninobu Yokota, *Professor*
Hideaki Kurata, *Associate Professor*
Tamotsu Yokota, *Associate Professor*
Shuichi Kato, *Assistant Professor*

General Summary

Physicians should aim patient-oriented medicine based on the concept of evidence-based medicine, which consists of research evidence, clinical expertise, and patients' preferences. To accomplish this goal, we encourage the members of our staff to do basic and clinical research. Areas of research include diabetes, metabolism, and endocrinology.

Research Activities

Epidemiology and evidence-based medicine

A nationwide epidemiologic study of mortality in approximately 3,500 patients with type 1 diabetes was started in 1986 and has continued to provide much information about the prognosis of Japanese children with type 1 diabetes. A population-based interventional study of childhood obesity and glucose intolerance has also continued. Several clinical trials of the treatment of type 2 diabetes using continuous glucose monitoring (CGM) are under way.

Genetics

Objective: Spontaneous hypoglycemia occurs owing to several causes with different patterns of hypoglycemia and hormone responsiveness. The aim of this study was to identify gene mutations in a family with spontaneous hypoglycemia by focusing on candidate genes and evaluating metabolism and hormone status.

Methods: The metabolic state was observed with CGM during the starvation test in the proband. Genomic DNA from peripheral blood was sequenced directly to identify gene mutations.

Results: The proband was a 34-year-old woman who was admitted to our university hospital because of severe hypoglycemia and metabolic acidosis associated with diarrhea and loss of appetite. She had had hypoglycemia-like episodes, especially when fasting, since the age of 1 year. In the starvation test, CGM clearly demonstrated no hypoglycemia until 29 hours. However, once hypoglycemia occurred at 29 hours, it persisted even after the induction of glucagon and the suppression of insulin secretion. These findings strongly suggest that a glyconeogenic enzyme is inactive. Therefore we focused on key glyconeogenic enzymes, including fructose-1,6-bisphatase (FBP1), phosphoenolpyruvate carboxykinase 1, and pyruvate kinase. The sequencing of these enzymes revealed that

the proband and her brother, who had similar hypoglycemia-like episodes, share the same mutant genotype of compound heterozygosity for *FBPI* (G164S/F194S), in which homozygotes of each allele had been reported as a responsible mutation for the phenotype.

Conclusion: Observation of hypoglycemia with CGM and hormone responsiveness in a patient with hypoglycemia permitted a focus on candidate genes and enabled identification of *FBPI* mutations.

Insulin resistance and obesity

A series of basic research studies of insulin resistance were performed in Otsuka Long-Evans Tokushima Fatty rats. The effects of a new oral hypoglycemic agent (a dipeptidyl peptidase IV inhibitor) on insulin resistance were investigated.

Dietary therapy

A highly monounsaturated enteral formula more effectively suppressed postprandial hyperglycemia without causing exaggerated insulin secretion compared with a high-carbohydrate enteral formula in patients with type 2 diabetes and healthy subjects. In patients with type 2 diabetes receiving tube feeding a highly monounsaturated enteral formula was shown with CGM to suppress postprandial hyperglycemia and to reduce 24-hour glycemic variations to greater extents compared with a high-carbohydrate enteral formula, even if carbohydrate nutrients had been adjusted for a low glycemic index.

Diabetic Vascular Complications

Diabetic complications are major sources of morbidity and mortality in patients with diabetes and an economic burden on societies worldwide. A greater understanding of the molecular targets that regulates both microangiopathy and macroangiopathy could lead to novel therapeutic strategies against diabetic complications. The Rho GTPases and their downstream effectors, Rho-associated kinases (ROCKs), have been implicated as regulators of the actin cytoskeleton. Because changes in the actin cytoskeleton are associated with vascular function, recent studies have revealed that ROCKs play a pivotal role in cardiovascular diseases, such as atherosclerosis, and in vascular remodeling. Accumulating evidence from animal models of diabetes shows that ROCK activity is increased in the kidney, retina, and vessels. Studies using pharmacological inhibition and genetic deletion of ROCKs have demonstrated that ROCK inhibition suppresses diabetic nephropathy by attenuating the excessive production of extracellular matrix induced by diabetes and slows the development of glomerular sclerosis and interstitial fibrosis. Given this background, we used db/db mice to study further mechanisms by which ROCKs regulate diabetic macroangiopathy. We found that ROCKs induce expression of monocyte chemoattractant protein 1 through activation of p38 mitogen-activated protein kinase and nuclear factor κ B under diabetic conditions. Furthermore, we found that ROCKs regulate the expression and function of hypoxia-inducible factor 1 α , thereby inducing glomerulosclerosis under diabetic conditions. Finally, we established a primary culture system for Schwann cells from diabetic mice.

Endocrinology

To identify and separate stem-like cells in human pituitary adenomas, we focused on the expression of CD133, which is a tumor stem cell marker in brain tumors, and examined the differences between CD133-positive cells and CD133-negative cells indicating stem properties.

The 12-lipoxygenase pathway may play a role in the pathogenesis of diabetic cardiomyopathy. Therefore, the role of the 12-lipoxygenase pathway in cardiomyopathy was examined in a rat model of diabetic cardiomyopathy and in an in-vitro study with a primary cardiomyocyte culture system.

Previous studies have shown that the secretion of adrenocorticotrophic hormone is increased in the hearts of patients with hypertension, indicating that adrenocorticotrophic hormone may be involved in the pathophysiology of cardiovascular diseases. Recently, pro-opiomelanocortin messenger RNA has been shown to be expressed in the murine heart. Therefore, we designed a study using HL-1 cardiomyocytes to clarify the pathophysiological role of pro-opiomelanocortin.

In patients with hyperaldosteronism, Ca blockers (type T and type N) reduce levels of aldosterone.

In patients with hypertension and type 2 diabetes, fluctuations of glucose and systolic blood pressure were found to be related.

Publications

Nakagami T, Nishimura R, Sone H, Tajima N. Accumulation of cardiovascular risks in Japanese women with abnormal glucose and mild to moderate hypercholesterolemia. *Int J Cardiol.* 2011; **152**: 254-6.

Nishimura R, Nakagami T, Sone H, Ohashi Y, Tajima N. Relationship between hemoglobin A1c and cardiovascular disease in mild-to-moderate hypercholesterolemic Japanese individuals: subanalysis of a large-scale randomized controlled trial. *Cardiovasc Diabetol.* 2011; **10**: 58.

Morimoto A, Nishimura R, Tsujino D, Taki K, Tajima N, Utsunomiya K. Relationship among A1C, hypoglycemia, and hyperglycemia in Japanese with type 2 diabetes—results from continuous glucose monitoring data. *Diabetes Technol Ther.* 2011; **13**: 667-70.

Tsujino D, Nishimura R, Taki K, Morimoto A, Tajima N, Utsunomiya K. Comparing the efficacy of α -glucosidase inhibitors in suppressing postprandial hyperglycemia using continuous glucose monitoring: a pilot study—the MAJOR study. *Diabetes Technol Ther.* 2011; **13**: 303-8.

Hiki Y, Sasaki T, Shimada K, Fujimoto K, Nemoto M, Utsunomiya K. Establishment of a novel transplantation method for murine islets embedded in reconstituted basement membrane matrix (in Japanese). *Tokyo Jikeikai Ika Daigaku Zasshi.* 2012; **127**: 49-61.

Shimada K, Tachibana T, Fujimoto K, Sasaki T, Okabe M. Temporal and spatial cellular distribu-

tion of neural crest derivatives and alpha cells during islet development. *Acta Histochem Cytochem.* 2012; **45**: 65-75.

Yamamoto M, Iguchi G, Takeno R, Okimura Y, Sano T, Takahashi M, Nishizawa H, Handyaningshi AE, Fukuoka H, Tobita M, Saitoh T, Tojo K, Mokubo A, Morinobu A, Iida K, Kaji H, Seino S, Chihara K, Takahashi Y. Adult combined GH, prolactin, and TSH deficiency associated with circulating PIT-1 antibody in humans. *J Clin Invest.* 2011; **121**: 113-9.

Nemoto M, Hiki Y, Shimada K, Nakai N, Fujimoto K, Inoue S, Sakurada N, Kaneko H, Sugita M, Okabe M, Sasaki T. Novel hormonal delivery method using the ink-jet technology: application to pulmonary insulin therapies. *Diabetes Technol Ther.* 2011; **13**: 509-17.

Ikeda K, Saito T, Tojo K. Efonidipine, a Ca(2+)-channel blocker, enhances the production of dehydroepiandrosterone sulfate in NCI-H295R human adrenocortical carcinoma cells. *Tohoku J Exp Med.* 2011; **224**: 263-71.

Kawanami D, Matoba K, Kanazawa Y, Ishizawa S, Yokota T, Utsunomiya K. Thrombin induces MCP-1 expression through Rho-kinase and subsequent p38 MAPK/NF- κ B signaling pathway activation in vascular endothelial cells. *Biochem Biophys Res Commun.* 2011; **411**: 798-803.

Mahabeshwar GH, Kawanami D, Sharma N, Takami Y, Zhou G, Shi H, Nayak L, Jeyaraj D,

Grealy R, White M, McManus R, Ryan T, Leahy P, Lin Z, Haldar SM, Atkins GB, Wong HR, Lingrel JB, Jain MK. The myeloid transcription factor KLF2 regulates the host response to polymicrobial infection and endotoxemic shock. *Immunity*. 2011; **34**: 715-28.

Hosoya T, Matsushima M, Nukariya K, Utsunomiya K. The relationship between the severity of depressive symptoms and diabetes-related emotional distress in patients with type 2 diabetes. *Intern Med*. 2012; **51**: 263-9.

Mori Y, Taniguchi Y, Sezaki K, Yokoyama J, Utsunomiya K. Liraglutide narrows the range of circadian glycemic variations in Japanese type 2 diabetes patients and nearly flattens these variations in drug-naïve type 2 diabetes patients: a continuous glucose monitoring-based study. *Diabetes Technol Ther*. 2011; **13**: 1139-44.

Mori Y, Tanaka T, Matsuura K, Yokoyama J, Utsunomiya K. Influence of telmisartan on insulin response after glucose loading in obese patients with hypertension: ARB trial of hypertension in

obese patients with hyperinsulinemia assessed by oral glucose tolerance test (ATHLETE). *Adv Ther*. 2011; **28**: 698-706.

Mori Y, Taniguchi Y, Matsuura K, Sezaki K, Yokoyama J, Utsunomiya K. Effects of sitagliptin on 24-h glycemic changes in Japanese patients with type 2 diabetes assessed using continuous glucose monitoring. *Diabetes Technol Ther*. 2011; **13**: 699-703.

Sumiyama K, Utsunomiya K, Ohya T, Aihara H, Ikeda K, Imazu H, Tamai N, Nagano H, Ishinoda Y, Tajiri H. A pilot study on ultrasound-assisted liposuction of the greater omentum in porcine models. *Minim Invasive Ther Allied Technol*. 2012; **21**: 71-7.

Mori Y, Shiozaki M, Matsuura K, Tanaka T, Yokoyama J, Utsunomiya K. Evaluation of efficacy of acarbose on glucose fluctuation and postprandial glucose using continuous glucose monitoring in type 2 diabetes mellitus. *Diabetes Technol Ther*. 2011; **13**: 467-70.